

After the Procedure

Contact the interventional radiologist's office **immediately** if you begin to bleed more than just oozing or you notice bright red blood!

If there is a small amount of blood on the band aids, there is no need to be alarmed. This is normal. You may remove the band-aid and replace it with a new one several hours after your procedure.

If there is no oozing, you may remove the band aid completely. Keep the sites clean and dry. You may notice tenderness or a small knot at the puncture sites. These will disappear over several weeks.

Report a temperature, pus or redness to your staff, these are signs of infection.

Notify your providers as soon as you notice your access has clotted, time is of the essence!



Heartland Kidney Network

7306 NW Tiffany Springs Pkwy
Suite 230
Kansas City, MO 64153

Phone: (816) 880-9990
Main Fax: (816) 880-9088
Data Fax: (816) 880-1775

Web: heartlandkidney.org
Email: net12@nw12.esrd.net



FISTULA COMPLICATIONS: STENOSIS & THROMBOSIS; NON-SURGICAL INTERVENTIONS



Promoting and facilitating high quality care standards for dialysis and kidney transplant patients in Iowa, Kansas, Missouri and Nebraska.

**PATIENT
EDUCATION
SERIES**



Complications Arise

Your body may not be used to the changes caused by your fistula. As a result, complications may arise and require medical attention.

INSIDE!

- ★ **Stenosis & Thrombosis Defined**
- ★ **Angiography & Radiology**
- ★ **After-Care & Follow-up**

This publication has been approved as appropriate community education by the Centers for Medicare & Medicaid Service's Fistula First Breakthrough Initiative Members. Feel free to copy this resource in its entirety for educational purposes. 2006.

COMPLICATIONS & INTERVENTIONS

Stenosis and Thrombosis



The continual force of blood rushing through your veins from your fistula can cause narrowing or stenosis, pronounced “sten-oh-sis” in your fistula. Another complication due to high blood flows is the formation of scar tissue inside your fistula. This scar tissue formation is sometimes referred to as “intimal hyperplasia”, pronounced “in-tim-all high-per-play-sha”. This condition can lead to decreased blood flows in your fistula or even clotting of your fistula from a blood clot.

If either occurs, dialysis may not be possible and your renal treatment team may schedule you an appointment with a doctor called an interventional radiologist, pronounced “in-ter-ven-shun-ul raid-eeaw-low-gist”. This doctor may perform angioplasty, pronounced “angee-oh-plas-tee”, or thrombolysis, pronounced “throm-bow-lie-sis” to allow your fistula to work correctly.

What is Interventional Radiology and Angiography?

Interventional radiology offers an alternative to surgery and has many patient benefits. Patients experience a reduction in pain and risks, a decrease in recovery times and costs as well as a reduction in hospitalizations. Most procedures can be done on an “outpatient” basis so no hospital stay is required.

Angiography, pronounced “an-gee-aw-gra-fee”, is an x-ray study of arteries. It is used to look at abnormalities in the blood vessels, such as narrowing or blockages, blood clots, and aneurysms, pronounced “an-your-is-m’s”. It helps the doctor determine the best treatment for you.

Angioplasty

Venous angioplasty is a procedure to open and preserve the working of a fistula due most typically to stenosis. During the procedure, the doctor guides a small wire through your fistula while using an x-ray machine to “see” the vein and wire while he works.

Continued From Previous Column

Once the doctor is able to “see” the area that is narrowed, he inserts a small tube over the wire. A tiny balloon is attached to the tube and advanced to the narrowed portion. The balloon is inflated to deflated to open the narrowing. The procedure is done through a needle puncture similar to your routine dialysis treatment. No hospital stay is usually required.

Thrombolysis

If your fistula becomes clotted or thrombosed, pronounced “throm-boast”, blood can no longer flow through it, thus making dialysis impossible. The doctor may be able to remove the clot and restore blood flow. Needles, similar to those used in dialysis are inserted and medicine to dissolve the clot is injected. Similarly, a device may be inserted to break up the clot & remove it from your access.

The doctor determines which technique is used. Usually a narrowed vein caused the clotting and you are usually then given venous angioplasty to treat the narrowed portion of your fistula.